



ecoinvent 3.11 Dataset Documentation

'power sawing, without catalytic converter - RER'


Note: This document contains only an extract of the information in the dataset. Additional data about properties of exchanges, mathematical relations, parameters, and contact information for authors and reviewers are available in the full dataset, i.e. in ecoSpold format. Amount and identity of the exchanges in an undefined dataset are independent of modeling choices of the different system models. Linked dataset are available in separate documents.

Table of Contents

Dataset Identification	3
Dataset Authorship	4
Exchange Summary	4
Dataset Description	5
Detailed Information For Exchanges	7
Reference product	7
Inputs from technosphere	7
Emissions to air	7
Emissions to soil	9
Source	10
Restriction of Use	10

Dataset Identification

Activity name	power sawing, without catalytic converter
Geography	Europe
Time period	2011-01-01 to 2012-12-31 - Valid for the entire period
Synonyms	chainsaw, chain saw
ISIC rev.4 ecoinvent	1610: Sawmilling and planing of wood
Reference product	power sawing, without catalytic converter
CPC classification	88311: Wood manufacturing services
Dataset type	Ordinary transforming activity
Technology level	Current
Version - system model	3.11 - Undefined



■ Geography ■ Water ■ Other

Dataset Authorship

Data generator	Frank Werner, Werner Umwelt & Entwicklung
Data entry	Frank Werner, Werner Umwelt & Entwicklung
Review	Emilia Moreno Ruiz, ecoinvent Centre

Exchange Summary

Reference product	Material for treatment	Byproduct classification	Amount
power sawing, without catalytic converter	None	allocatable product	1 hour

Inputs from technosphere	Amount
petrol, two-stroke blend	1.6 kg
power saw, without catalytic converter	0.0004 unit
vegetable oil, refined	0.54 kg

Emissions to air	Amount
Acetaldehyde	0.000687 kg
Acetone	0.000157 kg
Acrolein	4.41e-05 kg
Benzaldehyde	0.000461 kg
Benzo(a)pyrene	2.38e-07 kg
Carbon dioxide, fossil	2.15 kg
Carbon monoxide, fossil	0.981 kg
Dinitrogen monoxide	8.64e-05 kg
Formaldehyde	0.0351 kg
Methane, fossil	0.0413 kg
NM VOC, non-methane volatile organic compounds	0.298 kg
Nitrogen oxides	0.0086 kg
PAH, polycyclic aromatic hydrocarbons	5.95e-05 kg
Pentane	0.0342 kg
Propanal	0.000124 kg
Sulfur dioxide	3.2e-05 kg


Toluene	0.0332 kg
---------	-----------

Emissions to soil	Amount
-------------------	--------

Oils, non-fossil	0.054 kg
------------------	----------

Dataset Description

General comment



The inventory represents the operation of a professional power saw with a fuel consumption of 1.6 kg of standard two-stroke petrol blend (0.75 kg/l) for power saws per working hour and with a power output of about 3.1 - 3.2 kW.

Included activities start

Service beginning with the input of fuel into the power saw.

Included activities end

Includes the input of machinery infrastructure, the input of fuel, lubricants/greases as well as their disposal, and the emissions into air from fuel consumption. Wood is not included in the dataset.

Sampling procedure

Producer data and data from several scientific papers (emission measurements from different types of power saws, different fuels and additives).

Extrapolations

See geography.

Technology comment

The module represents average technology used in Europe around 2010

Geography comment

Data sourced from several European studies

Detailed Information For Exchanges

Reference product	Annual prod.vol.	Amount
power sawing, without catalytic converter	3.48e+9 hour	1 hour
Production volume: 3.48e+9 hour		
Production volume comment: 1,00,000 chainsaws with 2500 PMH, weighted by RER GDP/Global GDP		

Inputs from technosphere	Amount
petrol, two-stroke blend	1.6 kg
Comment: average consumption per working hour; reference for normalizing values from different sources	
Uncertainty distribution: lognormal; GSD2: 1.22; Pedigree matrix: [4, 5, 1, 1, 1]	
Source: Willared, J. (2011)	
power saw, without catalytic converter	0.0004 unit
Comment: calculated based on producer information	
Uncertainty distribution: lognormal; GSD2: 1.22; Pedigree matrix: [4, 5, 1, 1, 1]	
Source: Kellenberger D. (2007)	
vegetable oil, refined	0.54 kg
Comment: lubricating oil for chain	
Uncertainty distribution: lognormal; GSD2: 1.21; Pedigree matrix: [3, 5, 1, 1, 1]	
Source: Kellenberger D. (2007)	

Emissions to air	Subcompartment	Amount
Acetaldehyde	non-urban air or from high stacks	0.000687 kg
Comment: various sources: Magnussen et al (2002), Magnussen et al. (2000), Spielmann et al. 2007) as adequate		
Uncertainty distribution: lognormal; GSD2: 1.77; Pedigree matrix: [1, 5, 1, 1, 5]		
Acetone	non-urban air or from high stacks	0.000157 kg
Uncertainty distribution: lognormal; GSD2: 2.00; Pedigree matrix: [1, 5, 1, 1, 5]		
Acrolein	non-urban air or from high stacks	4.41e-05 kg
Uncertainty distribution: lognormal; GSD2: 2.05; Pedigree matrix: [1, 5, 5, 1, 5]		
Benzaldehyde	non-urban air or from high stacks	0.000461 kg
Uncertainty distribution: lognormal; GSD2: 2.05; Pedigree matrix: [1, 5, 5, 1, 5]		
Benzo(a)pyrene	non-urban air or from high stacks	2.38e-07 kg
Uncertainty distribution: lognormal; GSD2: 2.33; Pedigree matrix: [1, 5, 5, 1, 5]		
Carbon dioxide, fossil	non-urban air or from high stacks	2.15 kg
Comment: calculated based on a C content of petrol of 0.847 kg C/kg and 43% of carbon content emitted as CO2 (personal communication power saw producer)		
Uncertainty distribution: lognormal; GSD2: 1.21; Pedigree matrix: [1, 5, 3, 3, 1]		

Carbon monoxide, fossil	non-urban air or from high stacks	0.981 kg
Comment: calculated based on a C content of petrol of 0.8472 kg C/kg and 31% of carbon content emitted as CO (personal communication power saw producer)		
Uncertainty distribution: lognormal; GSD2: 1.58; Pedigree matrix: [2, 5, 1, 1, 1]		
Dinitrogen monoxide	non-urban air or from high stacks	8.64e-05 kg
Uncertainty distribution: lognormal; GSD2: 1.77; Pedigree matrix: [1, 5, 1, 1, 5]		
Formaldehyde	non-urban air or from high stacks	0.0351 kg
Comment: various sources: Magnussen et al (2002), Magnussen et al. (2000), Spielmann et al. 2007) as adequate		
Uncertainty distribution: lognormal; GSD2: 1.77; Pedigree matrix: [1, 5, 1, 1, 5]		
Methane, fossil	non-urban air or from high stacks	0.0413 kg
Uncertainty distribution: lognormal; GSD2: 1.77; Pedigree matrix: [1, 5, 1, 1, 5]		
NM VOC, non-methane volatile organic compounds	non-urban air or from high stacks	0.298 kg
Comment: amount of NM VOC emissions that are not inventoried as individual substances based on total NM VOC emissions of 26% of carbon content (personal communication by power saw producer) and based on a C content of petrol of 0.857 kg C/kg and excluding PAH, which are inventoried separately		
Uncertainty distribution: lognormal; GSD2: 1.78; Pedigree matrix: [3, 5, 1, 3, 5]		
Nitrogen oxides	non-urban air or from high stacks	0.0086 kg
Comment: personal communication of power chain producer		
Uncertainty distribution: lognormal; GSD2: 1.77; Pedigree matrix: [1, 5, 1, 1, 5]		
PAH, polycyclic aromatic hydrocarbons	non-urban air or from high stacks	5.95e-05 kg
Comment: amount of PAH that are not inventoried as individual substances as quantified based on Magnussen et al (2002) and Magnussen et al. (2000)		
Uncertainty distribution: lognormal; GSD2: 2.28; Pedigree matrix: [1, 5, 1, 1, 5]		
Pentane	non-urban air or from high stacks	0.0342 kg
Uncertainty distribution: lognormal; GSD2: 2.00; Pedigree matrix: [1, 5, 1, 1, 5]		
Propanal	non-urban air or from high stacks	0.000124 kg
Uncertainty distribution: lognormal; GSD2: 2.00; Pedigree matrix: [1, 5, 1, 1, 5]		
Sulfur dioxide	non-urban air or from high stacks	3.2e-05 kg
Comment: calculated based on an assumed S content of 0.001 % (by mass) as the threshold value for the Swiss tax on sulfur in Petrol		
Uncertainty distribution: lognormal; GSD2: 1.17; Pedigree matrix: [1, 1, 1, 1, 1]		
Toluene	non-urban air or from high stacks	0.0332 kg
Uncertainty distribution: lognormal; GSD2: 2.00; Pedigree matrix: [1, 5, 1, 1, 5]		

Emissions to soil	Subcompartment	Amount
Oils, non-fossil	forestry	0.054 kg

Comment: 10% of the oil from lubricating the chain

Uncertainty distribution: lognormal; **GSD2:** 1.58; **Pedigree matrix:** [1, 5, 3, 3, 1]

Source

First author	Willared, J.
Title	Personal written communication, Mr. Willaredt, Husqvarna AB
Year	2011
<hr/>	
First author	Magnusson R.
Additional author(s)	Nilsson C, Andersson K., Andersson B., Gieling R., Wiberg K., Östman C., Rannug U.
Title	Determination of chemical composition and mutagenicity in particles from chainsaw exhaust; experimental set-up, stability and results from two different fuels
Year	2010
Journal	Environmental Technology
Volume number	21
Issue number	7
<hr/>	
First author	Kellenberger D.
Additional author(s)	Althaus H.-J., Jungbluth N., Künniger T.
Title	Life Cycle Inventories of Building Products
Year	2007
Volume number	7

Restriction of Use

The restrictions of use stipulated in the EULA remain applicable for this pdf documentation.
Copyright ecoinvent Association, 2024